



## STARTING A PATIENT ON XYWAV

### PATIENT AUTHORIZATION

Patients can see if they are eligible for JazzCares services by completing the Patient Authorization Form at [jazzcares.com/xywav](http://jazzcares.com/xywav)

### NURSING SUPPORT TEAM

A dedicated team of nurses provides personalized support to your patients from the moment XYWAV is prescribed

### PHARMACY SUPPORT

Pharmacists at the Certified Pharmacy answer patients' questions about their XYWAV prescription 24/7



## COVERAGE AND REIMBURSEMENT

### DEDICATED SPECIALTY PHARMACY TEAM

Helps patients make the transition to XYWAV smoothly. Most patients transition to XYWAV in <30 days<sup>1</sup>

### PRIOR AUTHORIZATION (PA) AND APPEAL SUPPORT

Coverage details are often identified in <1 business day. If a PA is required, the Certified Pharmacy will provide your office with a unique CoverMyMeds<sup>®</sup> key code

### ANNUAL REVERIFICATION AND PA RECERTIFICATION

Prompts that help you complete a PA through CoverMyMeds in a timely manner



## FINANCIAL ASSISTANCE PROGRAMS

### QUICK START VOUCHER<sup>a</sup>

Eligible patients may be able to receive a **free 1-month supply** of XYWAV

### COUPON PROGRAM<sup>a</sup>

>99% of all eligible patients **can pay as little as \$5** for XYWAV<sup>2</sup>

### BRIDGE PROGRAM<sup>a</sup>

Helps prevent interruptions in therapy if insurance status changes. Eligible patients may **receive up to 120 days of free medication**

### PATIENT ASSISTANCE PROGRAM<sup>a</sup>

Eligible patients may get **access to free medication**. Visit [jazzcares.com/xywav](http://jazzcares.com/xywav) to complete and submit the application



## ONGOING SUPPORT

### NURSING AND PHARMACY TEAMS

Pharmacists, JazzCares Nurse Educators, and JazzCares Nurse Case Managers provide ongoing support to your patients

### myWAV™ APP

Optional online tool that can help patients get started and offer support throughout treatment

### ONLINE REFILL SCHEDULING

**Makes it convenient for your patients to schedule their next shipment via email and text reminders**

### XYWAV MENTOR PROGRAM

**Connects patients 1:1 with a peer who is also taking XYWAV**

### WEEKLY STATUS REPORTS

Helps you stay on top of your patients' progress with weekly data updates via email or fax

Encourage your patients to submit the JazzCares patient authorization at [jazzcares.com/xywav](http://jazzcares.com/xywav)

## Indications and Usage

XYWAV is indicated for the treatment of cataplexy or excessive daytime sleepiness (EDS) in patients 7 years of age and older with narcolepsy, and for the treatment of idiopathic hypersomnia (IH) in adults.

Please see additional Important Safety Information on next page and full [Prescribing Information](#), including **BOXED Warning**.

<sup>a</sup>Terms and conditions apply. Must be enrolled in the XYWAV and XYREM REMS. Jazz Pharmaceuticals, Inc. reserves the right to terminate or modify these programs at any time with or without notice. Other terms and conditions apply.

References: 1. Data on File-REF-8210. Jazz Pharmaceuticals, Inc. 2. Data on File-REF-8209. Jazz Pharmaceuticals, Inc.

## Important Safety Information

### WARNING: CENTRAL NERVOUS SYSTEM DEPRESSION and ABUSE AND MISUSE.

#### • **Central Nervous System Depression**

XYWAV is a CNS depressant. Clinically significant respiratory depression and obtundation may occur in patients treated with XYWAV at recommended doses. Many patients who received XYWAV during clinical trials in narcolepsy and idiopathic hypersomnia were receiving CNS stimulants.

#### • **Abuse and Misuse**

The active moiety of XYWAV is oxybate or gamma-hydroxybutyrate (GHB). Abuse or misuse of illicit GHB, either alone or in combination with other CNS depressants, is associated with CNS adverse reactions, including seizure, respiratory depression, decreases in the level of consciousness, coma, and death.

Because of the risks of CNS depression and abuse and misuse, XYWAV is available only through a restricted program under a Risk Evaluation and Mitigation Strategy (REMS) called the XYWAV and XYREM REMS.

### **Contraindications**

XYWAV is contraindicated in combination with sedative hypnotics or alcohol and in patients with succinic semialdehyde dehydrogenase deficiency.

### **Warnings and Precautions**

- **CNS Depression:** Use caution when considering the concurrent use with other CNS depressants. If concurrent use is required, consider dose reduction or discontinuation of one or more CNS depressants (including XYWAV). Consider interrupting XYWAV treatment if short-term opioid use is required. After first initiating treatment and until certain that XYWAV does not affect them adversely, caution patients against hazardous activities requiring complete mental alertness or motor coordination such as operating hazardous machinery, including automobiles or airplanes. Also caution patients against these hazardous activities for at least 6 hours after taking XYWAV. Patients should be queried about CNS depression-related events upon initiation of XYWAV therapy and periodically thereafter.
- **Abuse and Misuse:** XYWAV is a Schedule III controlled substance. The rapid onset of sedation, coupled with the amnestic features of GHB particularly when combined with alcohol, has proven to be dangerous for the voluntary and involuntary user (eg, assault victim).
- **Respiratory Depression and Sleep-Disordered Breathing:** XYWAV may impair respiratory drive, especially in patients with compromised respiratory function. In overdoses of oxybate and with illicit use of GHB, life-threatening respiratory depression has been reported. Increased apnea and reduced oxygenation may occur with XYWAV administration in adult and pediatric patients. A significant increase in the number of central apneas and clinically significant oxygen desaturation may occur in patients with obstructive sleep apnea treated with XYWAV. Prescribers should be aware that sleep-related breathing disorders tend to be more prevalent in obese patients, in men, in postmenopausal women not on hormone replacement therapy, and among patients with narcolepsy.
- **Depression and Suicidality:** In clinical trials in adult patients with narcolepsy and IH, depression and depressed mood were reported in patients treated with XYWAV. In most cases, no change in XYWAV treatment was required. Two suicides and two attempted suicides occurred in adult clinical trials with oxybate (same active moiety as XYWAV). One patient experienced suicidal ideation and two patients reported depression in a pediatric clinical trial with oxybate. **Monitor patients for the emergence of increased depressive symptoms and/or suicidality while taking XYWAV, which require careful and immediate evaluation.**
- **Other Behavioral or Psychiatric Adverse Reactions:** Monitor patients for impaired motor/cognitive function or the emergence of or increase in anxiety and/or confusion. The emergence or increase in the occurrence of behavioral or psychiatric events in patients taking XYWAV should be carefully monitored.
- **Parasomnias:** In clinical trials, parasomnias including sleepwalking were reported in adult patients treated with XYWAV. Parasomnias, including sleepwalking, also have been reported in a pediatric clinical trial with sodium oxybate (same active moiety as XYWAV) and in postmarketing experience with sodium oxybate. Episodes of sleepwalking should be fully evaluated and appropriate interventions considered.

### **Most Common Adverse Reactions**

The most common adverse reactions (occurring in  $\geq 5\%$  of XYWAV-treated patients in adult clinical trials in either narcolepsy or IH) were nausea, headache, dizziness, anxiety, insomnia, decreased appetite, hyperhidrosis, vomiting, diarrhea, dry mouth, parasomnia, somnolence, fatigue, and tremor.

In the pediatric clinical trial with XYREM (same active moiety as XYWAV) in patients 7 years of age and older with narcolepsy, the most common adverse reactions ( $\geq 5\%$ ) were nausea (20%), enuresis (19%), vomiting (18%), headache (17%), weight decreased (13%), decreased appetite (9%), dizziness (8%), and sleepwalking (6%). The safety profile in pediatric patients with XYWAV is expected to be similar to that of adult patients treated with XYWAV and to that of pediatric patients treated with XYREM.

Please see full **Prescribing Information**, including **BOXED Warning**.

For more information on XYWAV and JazzCares, visit [www.xywavhcp.com](http://www.xywavhcp.com) and [www.jazzcares.com/hcp/xywav](http://www.jazzcares.com/hcp/xywav).

XYWAV is a registered trademark of Jazz Pharmaceuticals, Inc.